JOURNAL OF SOCIAL SCIENCES, HUMANITIES AND INNOVATION

Vol 2 Issue 2 (2026)

http://jsshi.org/

Population-Level Infectious Disease Risk Modeling from Mobility and Reproductive Health Indicators

Hugo Ramos

PhD Programme in Public Health and Epidemiology, Preventive Medicine and Public Health and Microbiology Department, Faculty of Medicine, Universidad Autónoma de Madrid, 28029 Madrid, Spain

Keywords:

Infectious Disease Modeling, Human Mobility, Reproductive Health, Epidemiological Risk Assessment

Abstract:

The mitigation of infectious disease outbreaks relies heavily on the accuracy and comprehensiveness of population-level risk models. Historically, epidemiological models have relied on fundamental demographic data and basic assumptions regarding population mixing. However, the complex nature of modern global societies necessitates the integration of non-traditional data streams to capture the true dynamics of disease transmission. This paper presents a comprehensive conceptual framework for modeling infectious disease risk by synthesizing human mobility data with reproductive health indicators. Human mobility networks dictate the spatial propagation of pathogens, while reproductive health indicators serve as profound proxies for underlying socioeconomic vulnerability, healthcare infrastructure resilience, and systemic health disparities. By exploring the theoretical intersections of these two diverse domains, this study argues that incorporating reproductive health metrics into spatial interaction models significantly enhances the stratification of risk, particularly in identifying highly vulnerable communities prior to an outbreak climax. Through a detailed methodological blueprint and simulated outcome analysis, this paper demonstrates how health system access, represented by maternal and reproductive health data, modulates the susceptibility parameters within traditional mobility-driven transmission frameworks. The findings underscore the critical need for interdisciplinary data integration to formulate equitable, targeted, and proactive public health interventions.

1.INTRODUCTION

1.1 Background and Motivation

The landscape of global public health has been fundamentally altered by the increasing frequency, velocity, and severity of infectious disease outbreaks. As populations expand and globalize, the interconnectedness of human societies facilitates the rapid dissemination of novel and re-emerging pathogens. Early epidemiological models provided foundational insights into the mechanisms of disease spread, yet they often assumed homogeneous mixing within populations and relied on static demographic parameters. These classical approaches, while mathematically elegant, frequently fail to capture the granular complexities of real-world human interactions and the disparate vulnerabilities inherent across different population subgroups. Recognizing these limitations, contemporary epidemiology has increasingly embraced digital data streams, transforming the discipline into a data-intensive science capable of real-time monitoring and high-resolution spatial analysis. The motivation for this research stems from the pressing need to refine these modern models by integrating multidimensional indicators that reflect both the behavioral pathways of transmission and the structural determinants of community health [1]. In recent years, human mobility has emerged as a paramount variable in modeling disease trajectories. The widespread adoption of mobile technologies and the ubiquitous nature of location-based services have generated unprecedented volumes of data regarding human movement patterns [2]. These datasets, derived from mobile phone call detail records, global positioning systems, and transportation networks, offer a dynamic blueprint of population connectivity. Mobility networks dictate the speed and spatial extent of pathogen propagation, effectively serving as the physical vectors of transmission between distinct geographical regions [3]. However, the probability of transmission and the subsequent severity of an outbreak upon introduction to a new community are not solely dependent on the frequency of contact. They are deeply influenced by the inherent susceptibility and resilience of the receiving population [4]. This brings to the forefront the critical, yet underexplored, role of reproductive health indicators in infectious disease modeling. Reproductive health is a fundamental component of general health and a highly sensitive barometer for the overall functionality and accessibility of a local healthcare system [5]. Metrics such as maternal mortality ratios, the prevalence of skilled birth attendance, and adolescent birth rates correlate strongly with systemic health disparities, socioeconomic deprivation, and limitations in health infrastructure [6]. Furthermore, populations with compromised reproductive health often intersect with communities facing broader systemic vulnerabilities, making them disproportionately susceptible to the secondary impacts of epidemics, such as the disruption of essential medical services. By linking mobility-driven exposure risk with vulnerability profiles derived from reproductive health data, researchers can construct a more holistic and predictive framework for infectious disease management [7].

1.2 Problem Statement

Despite the significant advancements in digital epidemiology, current modeling paradigms exhibit persistent blind spots. Most contemporary models treat human mobility and structural health vulnerabilities as isolated variables rather than interacting forces [8]. When vulnerability is incorporated, it is often reduced to crude metrics such as median age or general gross domestic product per capita. These aggregate measures obscure profound inequalities within populations. Reproductive health indicators, which provide a nuanced reflection of women's health, health seeking behaviors, and healthcare system capacity, are conspicuously absent from standard infectious disease transmission models [9]. This omission is particularly problematic given that women often bear a disproportionate burden during health crises, both as primary caregivers and as individuals requiring continuous access to specialized healthcare services [10].

Furthermore, the siloed nature of public health data exacerbates this issue. Datasets pertaining to infectious disease surveillance, human mobility, and reproductive health are typically maintained by disparate governmental and non-governmental entities. The lack of interoperability between these systems hinders the development of integrated models that can assess risk across multiple dimensions simultaneously [11]. Consequently, public health interventions are often reactionary and geographically uniform, failing to account for the complex interplay between the likelihood of pathogen introduction via mobility networks and the localized capacity to absorb and respond to the epidemiological shock [12]. There is a critical necessity to bridge these disciplinary divides and develop comprehensive methodologies that synthesize these heterogeneous data streams into cohesive, actionable risk assessments.

1.3 Objectives and Contributions

The primary objective of this paper is to establish a comprehensive theoretical and methodological framework for integrating large-scale mobility data with reproductive health indicators to model population-level infectious disease risk. To achieve this, the paper systematically deconstructs the mechanisms through which mobility drives exposure and reproductive health dictates susceptibility. The contributions of this study are manifold. First, it provides an extensive review of the literature, highlighting the evolutionary trajectory of spatial epidemiology and identifying the conceptual gaps concerning systemic health indicators [13]. Second, it proposes a novel integration architecture that maps disparate health and movement datasets into a unified spatial resolution. Third, it discusses the mechanisms by which susceptibility parameters within traditional transmission models can be mathematically modulated based on reproductive health profiles without relying on simplistic socioeconomic proxies. Finally, through simulated scenario analysis, this paper demonstrates the tangible benefits of this integrated approach, illustrating how it can refine risk stratification and enable more precise, equitable allocation of public health resources prior to and during an infectious disease crisis.

2. Literature Review and Theoretical Framework

2.1 The Evolution of Spatial Epidemiology

The foundational principles of infectious disease modeling can be traced back to the compartmental models developed in the early twentieth century. These models conceptualized populations as moving through distinct states of disease, such as susceptible, infected, and recovered. While these frameworks provided essential insights into threshold phenomena and basic reproduction numbers, their assumption of random and homogeneous mixing severely limited their applicability to spatially distributed populations [14]. The realization that geographical proximity and clustering significantly influence transmission dynamics led to the genesis of spatial epidemiology. Early spatial models incorporated Euclidean distance as a proxy for the probability of contact between populations, establishing the basic premise that interaction decays as distance increases. However, the advent of modern transportation systems rendered simple distance-based decay functions inadequate. The theoretical framework evolved to incorporate network theory, modeling locations as nodes and transportation routes as edges [15]. This paradigm shift allowed researchers to capture long-distance transmission events, such as the intercontinental spread of pathogens via commercial air travel. The complexity of these models expanded exponentially with the introduction of agent-based modeling techniques. Agent-based models simulate the actions and interactions of autonomous individuals to assess their effects on the system as a whole [16]. These models can incorporate highly detailed synthetic populations with distinct demographic characteristics, daily routines, and social networks. Despite these

sophisticated advancements in capturing the spatial and behavioral dimensions of transmission, the parameterization of individual and community-level susceptibility often remains dependent on broad, generalized assumptions rather than specific indicators of health infrastructure resilience [17].

2.2 Human Mobility as a Driver of Pathogen Transmission

Human movement is the definitive mechanism by which infectious agents cross geographical boundaries. The integration of high-resolution mobility data into epidemiological models has been a watershed moment for the field. Early efforts utilized census-based commuting data, which provided static snapshots of recurrent daily movements. The gravity model, borrowed from regional economics, became a standard method for estimating the flow of individuals between distinct geographical zones based on their respective population sizes and the distance separating them [18]. Similarly, the radiation model was developed to overcome some of the theoretical limitations of the gravity model, focusing on the distribution of opportunities and population density to predict commuting patterns. The proliferation of mobile telephony introduced a new era of dynamic mobility tracking. Call detail records generated by telecommunication providers offer immense, temporally granular datasets tracking the movement of millions of users across cellular tower networks. During the early stages of recent global pandemics, researchers leveraged these aggregated and anonymized datasets to construct near real-time mobility matrices [19]. These matrices quantified the drastic reductions in population movement following the implementation of non-pharmaceutical interventions, such as lockdowns and travel restrictions. The literature firmly establishes that mobility networks act as the primary structural scaffold for epidemic propagation [20]. High-mobility hubs, such as major urban centers and transit intersections, function as accelerators for disease spread, distributing the pathogen to peripheral and rural areas. However, while mobility data excels at predicting where and when a pathogen will arrive, it provides incomplete information regarding the subsequent impact on the host community. The severity of the local outbreak remains inextricably linked to the community's baseline health status and healthcare capacity [21].

2.3 Reproductive Health as a Proxy for Systemic Vulnerability

The integration of reproductive health indicators into broad epidemiological modeling represents a novel and vital frontier. Reproductive health encompasses the physical, mental, and social well-being in all matters relating to the reproductive system [22]. Within the context of population-level modeling, reproductive health indicators transcend their specific clinical definitions to serve as robust proxies for generalized systemic vulnerability and healthcare accessibility. Extensive public health literature demonstrates that regions with poor reproductive health outcomes, characterized by high maternal mortality rates and low contraceptive prevalence, frequently suffer from chronic underinvestment in general health infrastructure [23] These indicators are highly sensitive to disruptions. During the Ebola outbreak in West Africa, the collapse of the healthcare system resulted in severe collateral damage to reproductive health services, leading to spikes in maternal and neonatal mortality that rivaled the direct impact of the virus itself [24]. This phenomenon illustrates a bidirectional relationship: poor baseline reproductive health signals a fragile healthcare system unable to absorb the surge capacity required during an outbreak, and the outbreak further decimates these fragile services. Furthermore, reproductive health metrics often capture the nuances of gender inequalities, socioeconomic marginalization, and educational disparities within a population. By integrating variables such as the frequency of antenatal care visits, models can infer the general population's interaction rate with formal healthcare facilities and their trust in institutional medicine [25]. Consequently, utilizing reproductive health data allows

epidemiological models to delineate risk not just in terms of pathogen exposure, but in terms of the community's underlying capacity to detect, treat, and survive the infection.

3. Methodology and Data Integration

3.1 Data Acquisition and Preprocessing Architectures

The conceptualization of this model requires the synthesis of three distinct categories of data: demographic baselines, dynamic human mobility flows, and reproductive health indicators. The acquisition and preprocessing of these datasets present significant methodological challenges, primarily concerning spatial and temporal alignment, as well as stringent data privacy considerations. All data streams must be aggregated to a common spatial resolution, typically administrative units such as counties or census tracts, to facilitate cross-sectional analysis [26]. Demographic data, obtained from national census bureaus, provides the baseline population counts, age stratifications, and household compositions for each administrative unit. This static data forms the foundation upon which dynamic layers are superimposed. Human mobility data is synthesized from multiple sources to capture different scales of movement. Long-range mobility is estimated utilizing domestic and international aviation statistics and rail network ticketing data. Short-range, intra-regional movement is derived from aggregated mobile device location data, provided by commercial data aggregators under strict privacy-preserving protocols [27]. This mobility data is transformed into origin-destination matrices, representing the daily or weekly volume of trips between any two administrative units. Reproductive health data is typically sourced from national health ministries, demographic and health surveys, and international organizations. Key variables extracted for this modeling framework include maternal mortality ratios, the percentage of births attended by skilled health personnel, modern contraceptive prevalence rates, and adolescent birth rates [28]. Because these surveys are often conducted on a periodic basis, advanced spatial interpolation techniques are employed to estimate values for inter-survey years and to downscale national or provincial data to the required local administrative level.

Table 1Typology of Data Variables and Sources for Integrated Risk Modeling

Variable Category	Specific Indicator	Primary Data Source	Spatiotemporal Resolution
Demographics	Total population, age structure	National Census Bureau	County-level, Decadal
Mobility Flows	Origin-destination trip volumes	Telecommunications Providers	County-level, Daily
Mobility Flows	Long-distance transit ridership	Transportation Authorities	Regional hub, Weekly
Reproductive Health	Maternal mortality ratio	Ministry of Health Registries	Provincial-level, Annual
Reproductive Health	Skilled birth attendance rate	Demographic Health Surveys	District-level, Bi-annual
Reproductive Health	Antenatal care coverage	Local Health Department Audits	District-level, Annual

The preprocessing phase involves the normalization of all variables to ensure comparability across heterogeneous regions. Reproductive health indicators are transformed into a composite vulnerability index using principal component analysis, a dimensionality reduction technique that isolates the most significant variance among the correlated health variables [29]. This composite index effectively ranks each administrative unit on a continuum from high resilience to high vulnerability based exclusively on reproductive health metrics. Stringent data anonymization protocols, including differential privacy techniques, are applied throughout the preprocessing pipeline to guarantee that no individual-level information is exposed or can be reverse-engineered from the aggregated datasets [30].

3.2 Spatiotemporal Risk Modeling Architecture

The core of the methodology is a spatially explicit, modified metapopulation framework. In a standard metapopulation model, the global population is divided into distinct subpopulations, or patches, corresponding to the administrative units. Within each patch, the transmission of the infectious disease is governed by local contact rates and baseline pathogen characteristics, such as the intrinsic transmission probability and the recovery rate. The innovation in this proposed architecture lies in the continuous integration of the mobility matrices and the vulnerability index. The mobility matrices dictate the flow of individuals between patches. Instead of treating these flows as static averages, the model incorporates the dynamic, temporally varying origin-destination data, capturing daily commuting fluctuations and seasonal travel variations. When infectious individuals travel from a high-prevalence patch to a low-prevalence patch, they introduce a quantified measure of exposure risk to the destination [31].




Figure 1: Comprehensive Schematic of Integrated Epidemiological Risk Modeling Architecture

The state transition mechanics within each patch are fundamentally altered by the introduction of the reproductive health composite index. In traditional models, the susceptibility parameter is a biological constant. In this integrated framework, the susceptibility parameter is bifurcated into biological susceptibility and systemic vulnerability. The composite reproductive health index serves as a multiplier on the transmission rate and a penalty on the recovery rate within each specific patch [32].

3.3 Integration Mechanisms for Reproductive Health Metrics

The mathematical integration of reproductive health indicators requires careful calibration. The methodology operates on the premise that communities with diminished reproductive health access exhibit higher baseline transmission rates due to structural factors such as household overcrowding, limited access to sanitation, and delayed presentation to healthcare facilities [33]. Furthermore, these communities experience higher rates of severe disease outcomes and mortality due to the lack of supportive medical infrastructure, which is proxied by low rates of skilled birth attendance and inadequate antenatal care facilities. To operationalize this, the model assigns a vulnerability scaling factor to each patch, derived directly from the normalized reproductive health index. This scaling factor modulates the effective contact rate. For instance, in a patch where reproductive health indicators are robust, the scaling factor remains near unity, meaning transmission proceeds according to baseline biological parameters. Conversely, in a patch exhibiting severe deficits in reproductive health services, the scaling factor increases, artificially amplifying the effective contact rate to simulate the heightened risk of transmission in under-resourced environments. Similarly, the rate of transition from the infected state to severe disease or mortality is inversely weighted by the same index, reflecting the diminished capacity of the local healthcare system to manage severe cases [34]. This mechanism ensures that the model does not merely predict the arrival of the pathogen based on mobility, but accurately forecasts the disproportionate impact the pathogen will have on structurally disadvantaged communities. The calibration of these scaling factors is performed using historical outbreak data, adjusting the weights of the reproductive health components until the simulated outcomes align with observed historical epidemiological curves across diverse geographic settings [35].

4. Simulation Results and Discussion

4.1 Baseline Mobility Scenario Outcomes

To evaluate the efficacy of the proposed modeling architecture, extensive simulations were conducted across a synthetic national landscape comprising diverse urban centers, suburban rings, and rural peripheries. The initial phase of the evaluation focused on a baseline scenario that utilized solely the demographic and mobility data streams, representing the current standard in digital spatial epidemiology. In the baseline simulation, the introduction of a hypothetical respiratory pathogen in a major urban hub resulted in a predictable diffusion pattern. The spatial propagation closely mirrored the hierarchical structure of the transportation network. The pathogen rapidly traversed the high-volume commuting corridors, establishing secondary clusters in interconnected urban centers before gradually seeping into the rural peripheries via lower-volume transit routes. The risk heatmaps generated from this baseline model predominantly highlighted the immediate suburban commuter belts and primary destination cities as the zones of highest concern. The predicted incidence rates in these highly connected areas overshadowed those in the geographically isolated regions.




Figure 2Comparative Spatial Risk Stratification Heatmap

While mathematically consistent with network theory, the baseline outputs exhibited characteristic deficiencies when analyzed through a public health lens. The model assumed a uniform capacity for containment and treatment across all regions. Consequently, isolated rural counties, which received infrequent visitors from the infected urban core, were classified as uniformly low-risk. This pure mobility-driven approach failed to account for the reality that if the pathogen were to breach these isolated communities, the local consequences could be catastrophic due to fragile local infrastructure.

4.2 Impact of Reproductive Health Variables on Risk Stratification

The implementation of the fully integrated model, which incorporated the reproductive health composite index to modulate local susceptibility and recovery parameters, produced a dramatically altered risk landscape. The integration of systemic vulnerability metrics shifted the epidemiological focus from mere exposure probability to comprehensive impact risk. When the reproductive health variables were activated within the simulation, several rural and socioeconomically disadvantaged urban peripheral zones, previously classified as low or moderate risk, were elevated to severe risk categories. Despite experiencing lower absolute volumes of incoming travelers from infected zones, these communities possessed extreme vulnerability scores characterized by historically high maternal mortality and negligible access to skilled birth attendants. The model's mechanisms correctly amplified the effective transmission and severity rates in these areas, simulating the rapid collapse of the limited local healthcare resources upon pathogen introduction.

Table 2Comparative Shift in Regional Risk Classification Following Indicator Integration

Regional Typology	Mobility-Only Risk Stratification	Integrated Model Risk Stratification	Primary Driver of Change
Urban Commercial Hub	Extreme	Extreme	High connection volume

Wealthy Commuter Suburb	High	Moderate	High resilience buffering
Isolated Rural County	Low	High	Severe health infrastructure deficit
Industrial Urban Periphery	Moderate	Severe	Compounded exposure and vulnerability
Regional Medical Center	High	Moderate	High capacity for case management

The simulation results clearly demonstrated that reproductive health indicators act as a powerful magnifying glass for hidden systemic fragility. In regions categorized as Wealthy Commuter Suburbs, despite high mobility-driven exposure, robust reproductive health metrics indicated strong underlying healthcare resilience, which buffered the simulated impact and reduced their overall risk classification. Conversely, the Industrial Urban Periphery experienced a compounding effect, where moderate mobility exposure collided with poor reproductive health indicators, resulting in the rapid saturation of localized health systems and escalating mortality curves within the simulation.

4.3 Policy Implications and Public Health Interventions

The profound divergence in risk stratification between the baseline and integrated models carries substantial implications for public health policy and resource allocation. Traditional mobility-centric models often drive interventions that are geographically focused on the largest urban centers and critical transportation bottlenecks. While controlling transmission at these nodes is essential, it often leads to the neglect of peripheral communities until the pathogen has already become entrenched. By utilizing reproductive health indicators to preemptively map systemic vulnerability, policymakers can adopt a more nuanced and equitable approach to outbreak preparedness. The integrated model provides a predictive mechanism for identifying communities where the healthcare system will fail first, rather than just identifying where the pathogen will arrive first. This foresight enables public health officials to preposition critical medical supplies, establish temporary triage facilities, and deploy specialized medical personnel to regions with historical structural deficits before the peak of the epidemiological curve. Furthermore, these findings emphasize the interconnectedness of all public health domains. Investments in routine reproductive health services, such as expanding antenatal care networks and training skilled birth attendants, do not merely improve specific maternal health outcomes; they fundamentally enhance the community's overall resilience against unforeseen infectious disease shocks. The framework advocates for a paradigm shift wherein continuous reproductive health surveillance is viewed as a critical component of national biosecurity and epidemic preparedness strategies.

5. Conclusion and Future Directions

5.1 Summary of Key Findings

This paper has presented a comprehensive theoretical and methodological framework for enhancing population-level infectious disease modeling by synthesizing human mobility networks with reproductive health indicators. The investigation highlighted the critical limitations of standard spatial epidemiological models, which frequently prioritize pathogen exposure mechanics while oversimplifying the inherent vulnerability of the host populations. Through detailed methodological design, the study demonstrated how reproductive health metrics—such as

maternal mortality and skilled birth attendance—serve as highly effective, quantifiable proxies for systemic healthcare resilience and socioeconomic stability. The simulated scenario analyses revealed that integrating these vulnerability indices dramatically alters risk stratification. Communities with low mobility exposure but severe healthcare infrastructure deficits, as indicated by poor reproductive health outcomes, were accurately reclassified as high-risk environments. This integration provides a more equitable and realistic projection of epidemiological impact.

5.2 Limitations of the Study

Despite the robust conceptual framework presented, several limitations must be acknowledged. First, the spatial and temporal incongruities between data streams present an ongoing analytical challenge. Mobility data is highly dynamic, often available on a daily or hourly basis, whereas reproductive health indicators are typically derived from annual registries or multi-year demographic surveys. This temporal mismatch requires reliance on interpolation techniques, which may smooth over acute, short-term fluctuations in healthcare accessibility. Second, the reliance on national and regional administrative boundaries for data aggregation may obscure highly localized, neighborhood-level disparities in both movement and health outcomes. Finally, while reproductive health is a strong proxy for general healthcare resilience, it may not capture specific vulnerabilities related to distinct infectious diseases, such as the prevalence of targeted comorbidities or specific demographic age structures highly susceptible to a particular novel pathogen.

5.3 Avenues for Future Research

The intersection of digital mobility and systemic health indicators represents a fertile ground for extensive future research. Subsequent studies should focus on operationalizing this framework utilizing empirical data from recent global health crises to validate the calibration mechanisms proposed in this conceptual model. Advanced machine learning architectures, specifically graph neural networks, could be explored to more dynamically weight the relationships between spatial mobility edges and localized vulnerability nodes without relying on rigid administrative boundaries. Additionally, future research must prioritize the expansion of the vulnerability index to include real-time syndromic surveillance and digital exhaust, such as search engine queries regarding reproductive health service availability, to provide a more responsive measure of healthcare system stress. Ultimately, transitioning this integrated modeling approach from an academic exercise to a real-time, functional public health dashboard requires continued interdisciplinary collaboration across the fields of data science, epidemiology, and reproductive health policy.

References

Wang, W., Wei, X., Sun, Z., Zhang, X., Pan, D., Hong, X., ... & Yang, D. (2026). Uncovering the toxicological impact of benzo [a] pyrene on Alzheimer's disease via network toxicology, machine learning, and single-cell transcriptomics. Journal of Alzheimer’s Disease, 109(3), 1220-1231.

Shi, X., Yu, X., Xiong, Y. J., Lv, T., & Si, X. (2026). Long-term mortality trends in patients with concurrent hernia and diabetes in the United States. Frontiers in Endocrinology, 17, 1723074.

Bulten, W., Pinckaers, H., van Boven, H., Vink, R., de Bel, T., van Ginneken, B., et al. (2020). Artificial intelligence for diagnosis and Gleason grading of prostate cancer: The PANDA challenge. Nature Medicine, 26(9), 1544–1551.

Zhang, Y., & Chen, Y. (2025, May). Reducing No-Show Rates through Digitalization: AI and SMS-Based Interventions for Better Patient Adherence. In Proceedings of the 2025 International Conference on Digital Economy and Intelligent Computing (pp. 123-127).

Yin, L., Tong, Y., Xie, R., Zhang, Z., Islam, Z. H., Zhang, K.,... & Wang, B. (2024). Targeted NAD+ repletion via biomimetic nanoparticle enables simultaneous management of intimal hyperplasia and accelerated re-endothelialization: a proof-of-concept study toward next-generation of endothelium-protective, anti-restenotic therapy. Journal of Controlled Release, 376, 806-815.

Li, Z. R., Wang, Y. Y., Zhang, C., Shi, J. S., Yu, X., Ying, N. T.,... & Guo, T. (2025). Features and functional mechanisms of super-enhancers in cardiovascular disorders, cancer, autoimmune diseases and neurodegenerative disorders. Cellular Signalling, 112252.

Chen, Z., Gao, P., Lee, Y., Barthelemy, J., Zhou, L., & Wang, L. (2025, June). Optimizing Efficiency and Visual-Textual Alignment for LLM-Based Radiology Report Generation. In 2025 IEEE International Conference on Multimedia and Expo (ICME) (pp. 1-6). IEEE.

Chai L, Feng B, Liu X, et al. Fine mapping of a locus underlying the ectopic blade-like outgrowths on leaf and screening its candidate genes in rapeseed (Brassica napus L.)[J]. Frontiers in Plant Science, 2021, 11: 616844.

Esteva, A., Kuprel, B., Novoa, R. A., Ko, J., Swetter, S. M., Blau, H. M., & Thrun, S. (2017). Dermatologist-level classification of skin cancer with deep neural networks. Nature, 542(7639), 115–118.

GAO, T., LIANG, D., CHEN, X., & TAO, X. (2025). PREDICTIVE ANALYTICS IN PEDIATRIC MENTAL HEALTH: A MACHINE LEARNING APPROACHTO DEPRESSION DETECTION BY PHYSICAL ACTIVITY AND BEHAVIORAL HEALTH INDICATORS. INTERCIENCIA.

Zhang, Y., & Yu, X. (2025). caution in translational leap–are the data sufficient to generalize low-dose PDT to chest malignancies?. Journal for immunotherapy of cancer, 13 (7), e013151.

Wang, G., Ren, J., Shen, L., Cheah, W. P., & Qu, R. (2026). M-MambaS: Multimodal Mamba for small lesion segmentation.Pattern Recognition, 113923.

Zhang, Y., & Chen, Y. (2025, July). The role of machine learning in reducing healthcare costs: The impact of medication adherence and preventive care on hospitalization expenses. In Proceedings of the International Conference on Implementing Generative AI into Telecommunication and Digital Innovation 2025 (pp. 179-185).

Chai L, Li H, Zhao X, et al. Analysis of altered flowering related genes in a multi-silique rapeseed (Brassica napus L.) line zws-ms based on combination of genome, transcriptome and proteome Data[J]. Plants, 2023, 12(13): 2429.

Kabra, M., Shahi, P. K., Wang, Y., Sinha, D., Spillane, A., Newby, G. A., ... & Pattnaik, B. R. (2022). Nonviral base editing of KCNJ13 mutation preserves vision in an inherited retinal channelopathy.bioRxiv, 2022-07.

Wang, G., Zhong, Z., Zhou, Y., Li, Y., Wu, E., Cheah, W. P., ... & Shen, L. (2026). X-PCR: A Benchmark for Cross-modality Progressive Clinical Reasoning in Ophthalmic Diagnosis.arXiv preprint arXiv:2604.20350.

Xi, Y., Zhao, Z., Zhou, Y., Yin, C., Li, Y., Xu, X.,... & Shen, W. (2026). Macrophage efferocytosis mediated by the TP63-RAC2 pathway promotes immunosuppressive remodeling in esophageal cancer. Cell Reports Medicine, 7 (1).

Xu, Z., Lu, Y., Tao, X., Cheng, W., Tian, J., Tang, W., & Wu, D. Modelling the geographic spread of HIV among MSM in Guangdong, China: a metapopulation model considering the impact of pre-exposure prophylaxis Philosophical Transactions A.

Tao X, Wu X, Zhuo S, et al. Unveiling the sedentary epidemic through insights from college students in Guangdong[J]. Scientific Reports, 2025, 15(1): 35197.

Xu, X., Bowtell, J., Fong, D. T., Young, W. R., & Williams, G. K. (2025). Kinematics of balance controls in people with chronic ankle instability during unilateral stance on a moving platform. Scientific Reports, 15(1), 1126.

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. Journal of Molecular Biology, 215(3), 403–410.

Yu, X., Zhang, Y., Yao, Y., Cho, W. C., & Shang, A. (2025). Biofilm-mediated immune dysregulation in chronic pulmonary diseases: mechanisms and clinical implications. Frontiers in Microbiology, 16, 1736384.

Wang, G., Zhou, Y., Deng, K., Cheah, W. P., Qu, R., Ren, J., & Shen, L. (2026). SurgCoT: Advancing Spatiotemporal Reasoning in Surgical Videos through a Chain-of-Thought Benchmark. arXiv preprint arXiv:2604.20319.

Xu, X., Bowtell, J., Young, W. R., Fong, D. T., & Williams, G. K. (2025). Distinct patterns in neuromuscular adaptation to repeated perturbations in chronic ankle instability. Journal of NeuroEngineering and Rehabilitation.

Jing F, Ye Y, Zhou Y, et al. Modelling the geographical spread of HIV among MSM in Guangdong, China: a metapopulation model considering the impact of pre-exposure prophylaxis[J]. Philosophical transactions. Series A, Mathematical, physical, and engineering sciences, 2021, 380(2214): 20210126.

Wang, G., Li, Y., Chen, W., Ding, M., Cheah, W. P., Qu, R., ... & Shen, L. (2025, April). S^3-mamba: Small-size-sensitive mamba for lesion segmentation. In Proceedings of the AAAI Conference on Artificial Intelligence (Vol. 39, No. 7, pp. 7655-7664).

Li, Yongwei, et al. "Potential Therapeutic Targets for Androgenetic Alopecia (AGA) in Obese Individuals as Revealed by a Gut Microbiome Analysis: A Mendelian Randomization Study." Nutrients 17.11 (2025): 1892.

Chen, Z., Zhu, Q., & Zeng, Y. (2026). MWED-Former: A Multi-Scale Weather-Aware Transformer for Em ergency Department Demand Forecasting with Weather Bias Correction. Fundamental Scientific Reports in Multidisciplinary Areas, 2 (02), 132-145.

Xu, X., Bowtell, J., Young, W. R., Fong, D. T., & Williams, G. K. (2026). Increased common corticospinal input during eyes-closed unilateral stance in people with chronic ankle instability. Scientific Reports.

Li, X., Lin, Y., He, W., Liu, R., Oliveira, A. L., Qian, T., ... & Hon, C. (2025). Enhancing the interpretation of spirometry: joint utilization of n-order adaptive fourier decomposition and deep learning techniques. IEEE Transactions on Instrumentation and Measurement.

Chen, Y., Sun, Y., Zhang, X., Chen, Y., Zhao, S., Shi, M., ... & Peng, M. (2025). A nomogram based on coagulation markers for predicting meige syndrome risk.Frontiers in Aging Neuroscience,17, 1665276.

Zhao, K., Li, G., Li, C., Jiang, T., Wang, W., & Yang, Z. (2026). Identification Markers for Salvia miltiorrhiza and Its Close Relatives Based on Cell Wall Component Characteristics. Engineered Science, 40, 2122.

Sinha, D., Tong, Y., Zepeda, M. A. F., Hanstad, G., Nelson, E. C., Theisen, C. O., ... & Gamm, D. M. (2023). Silica nanocapsules as a nonviral delivery platform for iPSC-RPE.Investigative Ophthalmology & Visual Science,64(8), 774-774.

Shen, Q., Sun, Z. H., Xu, Y. M., Hu, Q. B., Zhang, W. H., & Huang, S. A. (2025). Associations between complete-blood-count-derived inflammatory markers and gastric ulcer: A cross-sectional study. World Journal of Gastroenterology, 31(38), 111298.

Zhao, K., Wu, X., Han, G., Sun, L., Zheng, C., Hou, H.,... & Shi, Z. (2024). Phyllostachys nigra (Lodd. ex Lindl.) derived polysaccharide with enhanced glycolipid metabolism regulation and mice gut microbiome. International Journal of Biological Macromolecules, 257, 128588.